

Fairbanks North Star Borough Standing Orders January 11, 2017

ERRATA Sheet

Key

Yellow – In previous versions

Blue – Corrections

Ketamine- Page 174

- o Duration: 4-12 Hours changed to Duration: 10-20 min

Dosage and Administration

• Adult

Behavioral Emergencies

- o 4 mg/kg IM

Excited Delirium

- o 4 mg/kg IM, (repeat in 5-10 min as needed) Max dose 500 mg. (First line Treatment for Excited Delirium is consider cause)

Dosage and Administration

• Adult

Behavioral Emergencies & Excited Delirium

- o 4 mg/kg IM, (repeat in 5-10 min as needed) Max dose 500 mg. (First line Treatment for Excited Delirium is consider cause)

Chest Pain- Page 36

Added the following to the Focused Exam: 15-lead if right heart MI or failure a possibility. In NSTEMI, with chest pain, always consider a 15-lead to rule out right heart or posterior MI.

Respiratory Distress: CHF/Pulmonary Edema- Adult- Page 86

Nitroglycerin (Nitrolingual):

- o Adult: 0.4 mg SL every 3-5 minutes PRN. If 3 doses ineffective, try CPAP

Nitroglycerin (Nitrolingual):

- o Adult: 0.4 mg SL every 3-5 minutes PRN. Use in conjunction with CPAP.

Cardiac: Tachycardia- Adult

Adenosine Triphosphate (Adenocard)

- 6 mg RAPID IV push, followed immediately by a 10-20 ml NS flush; If no response in 1-2 minutes, give 12 mg rapid IV push followed by a 10-20 ml NS flush. May give third dose of 12 mg rapid IV push followed by a 10-20 ml NS flush, if necessary

The remaining changes were spelling and typographical error.